

Dosage Chart



Children’s Motrin (Ibuprofen) Dosages:

Weight	Age	Infant’s Ibuprofen Concentrated Drops (50mg/1.25mL) Dropperful	Children’s Ibuprofen Suspension (100mg/5mL) mL (mililiter)	Children’s Soft Chewable Tablets (50mg) Tablets	Junior Strength Chewable Tablets (100mg) Tablets	Adult Tablets (200mg) Tablets
	0-5 mos.	Call Office	Call Office	Do Not Use	Do Not Use	Do Not Use
12-17 lbs.	6-11 mos.	1.25 mL	2.5 mL	Do Not Use	Do Not Use	Do Not Use
18-23 lbs.	12-23 mos.	1.875 ml	3.75 mL	Do Not Use	Do Not Use	Do Not Use
24-35 lbs.	2-3 yrs.		5 mL	2 tablets	Do Not Use	Do Not Use
36-47 lbs.	4-5 yrs.		7.5 mL	3 tablets	Do Not Use	Do Not Use
48-59 lbs.	6-8 yrs.		10 mL	4 tablets	2 tablets	1 tablet
60-71 lbs.	9-10 yrs.		12.5 mL	5 tablets	2.5 tablets	1 tablet
72-95 lbs.	11 yrs.		15 mL	6 tablets	3 tablets	1.5 tablets

NOTE: Not to be used under 6 months of age. One dose lasts 6-8 hours.
 Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp).
 Don’t use household teaspoons, which can vary in size.
 Aspirin should not be used in children to treat fever or pain.